

Polarization in interferometry

Michiel Brentjens

ASTRON, Dwingeloo, The Netherlands

NRAO Synthesis Imaging Workshop 2024-05-16

Reading list

- Born & Wolf *Principles of optics*
- Thompson, Moran & Swenson *Interferometry and Synthesis in Radio Astronomy*
- Taylor, Carilli & Perley *Synthesis Imaging in Radio Astronomy II*
- Bracewell *The Fourier Transform & Its Applications*
- Hamaker, Bregman & Sault *Understanding radio polarimetry: paper I*(1996)
- Sault, Hamaker & Bregman *paper II*(1996)
- Hamaker & Bregman *paper III* (1996)
- Hamaker *paper IV* (2000)
- Hamaker *paper V* (2006)
- Brentjens & de Bruyn *Faraday rotation measure synthesis* (2005)

Outline

EM wave physics


Astrophysics

Polarized EM-waves

Interferometric polarimetry


Messy reality

Electromagnetic (EM) wave




- **Vector** phenomenon
- From Maxwell's equations:
 $\hat{\mathbf{k}} = \hat{\mathbf{E}} \times \hat{\mathbf{B}}$
- We know $\mathbf{k}$
- Measure either $\mathbf{E}$ or $\mathbf{B}$
- $\mathbf{E}$ is easier

Electromagnetic (EM) wave



- **Vector** phenomenon
- From Maxwell's equations:
 $\hat{\mathbf{k}} = \hat{\mathbf{E}} \times \hat{\mathbf{B}}$
- We know $\mathbf{k}$
- Measure either $\mathbf{E}$ or $\mathbf{B}$
- $\mathbf{E}$ is easier

Electro-magnetic wave



- **Vector** phenomenon
- From Maxwell's equations:
 $\hat{\mathbf{k}} = \hat{\mathbf{E}} \times \hat{\mathbf{B}}$
- We know $\mathbf{k}$
- Measure either $\mathbf{E}$ or $\mathbf{B}$
- $\mathbf{E}$ is easier
- But:
- E_x and E_y **not equal**
- $\mathbf{E}$ may **rotate** as function of $\mathbf{x}$ and t .
- $\mathbf{E}$ traces **ellipse**

“Polarization”

Outline

EM wave physics


Astrophysics

Polarized EM-waves

Interferometric polarimetry

Messy reality

Synchrotron emission



Process

- Generates polarized emission
- Main emission mechanism at cm-m wavelength
- Up to 80% linearly polarized
- No circular
- $\langle \mathbf{E}_{\text{source}} \rangle \perp \mathbf{B}_{\text{source}}$



Polarimetry provides

- **B**-field direction
- Turbulence
- Indirectly: **B**-field strength



Zeeman splitting

H_I hyperfine transition



Process

- Generates polarized emission
- Only in spectral lines
- If magnetic moment: e.g. H_I , OH, CN, H_2O
- **B**-field splits RCP and LCP
- Separation: 2.8 Hz mG^{-1}

Polarimetry provides

- **B**-field strength at source
- If detectable...

Zeeman effect *Vlemmings et al. (2001)*






Fig. 4. Total power (*I*) and circular polarization (*V*) spectrum of the brightest H₂O maser feature around S Per. The dashed line is the fit of the synthetic *V*-spectrum to the observed spectrum. Also shown are the observed (dashed) and expected (solid) positions of the minimum and maximum of the *V*-spectrum.

Scattering/reflection

Thomson scattering




Process

- Modifies polarization state
- Thomson scattering: no T dependence
- Planets / Moon: dielectric transition


Polarimetry provides

- Electron densities in cool gas
- Dust properties
- Lunar dielectric constant

Faraday rotation



λ^2 law *Haverkorn et al. (2001)*



Process

- Modifies polarization state
- Delay between LCP and RCP
- Rotates linear pol angle
- $\Delta\chi = \chi_0 + \phi\lambda^2$

$$\phi = 0.812 \int_{\text{there}}^{\text{here}} n_e \mathbf{B} \cdot d\mathbf{l}$$

Polarimetry provides

- Source plasma properties
- Intervening plasma properties
- Rare cases: 3D tomography

ASTRON

Netherlands Institute for Radio Astronomy

Outline

EM wave physics

Astrophysics


Polarized EM-waves

Interferometric polarimetry

Messy reality

Polarization ellipse: cartesian basis

Geometry



Viewing from antenna towards source, watching orientation and length of $\mathbf{E}$ vector on a plane at a fixed location in space.


$$\mathbf{E} = E_x \hat{\mathbf{e}}_x + E_y \hat{\mathbf{e}}_y$$

$$E_x = A_x \cos(2\pi\nu t + \delta_x)$$


$$E_y = A_y \cos(2\pi\nu t + \delta_y)$$

- A_x = x-amplitude
- A_y = y-amplitude
- $\delta_{xy} = \delta_y - \delta_x$
- δ_{xy} = measure of ellipticity
- $\delta_{xy} > 0$: CW rotation $\Rightarrow$ LEP
- $\delta_{xy} = 0$: linear polarization
- $\delta_{xy} < 0$: CCW rotation $\Rightarrow$ REP


Polarization ellipse: xy 000




Polarization ellipse: xy 001




Polarization ellipse: xy 002




Polarization ellipse: xy 003




Polarization ellipse: xy 004




Polarization ellipse: xy 005




Polarization ellipse: xy 006




Polarization ellipse: xy 007




Polarization ellipse: xy 008




Polarization ellipse: xy 009




Polarization ellipse: xy 010




Polarization ellipse: xy 011




Polarization ellipse: xy 012




Polarization ellipse: xy 013




Polarization ellipse: xy 014




Polarization ellipse: xy 015




Polarization ellipse: xy 016




Polarization ellipse: xy 017




Polarization ellipse: xy 018




Polarization ellipse: xy 019




Polarization ellipse: xy 020




Polarization ellipse: xy 021




Polarization ellipse: xy 022




Polarization ellipse: xy 023




Polarization ellipse: xy 024




Polarization ellipse: xy 025




Polarization ellipse: xy 026




Polarization ellipse: xy 027




Polarization ellipse: xy 028



Polarization ellipse: xy 029




Polarization ellipse: xy 030



Polarization ellipse: circular basis

Geometry



Viewing from antenna towards source, watching orientation and length of $\mathbf{E}$ vector on a plane at a fixed location in space.


$$\mathbf{E} = A_r \hat{\mathbf{e}}_r + A_l \hat{\mathbf{e}}_l$$

$$\hat{\mathbf{e}}_r = \begin{pmatrix} \cos(2\pi\nu t + \delta_r) \\ \sin(2\pi\nu t + \delta_r) \end{pmatrix}$$


$$\hat{\mathbf{e}}_l = \begin{pmatrix} \cos(2\pi\nu t + \delta_l) \\ -\sin(2\pi\nu t + \delta_l) \end{pmatrix}$$

- $A_r + A_l =$ semi-major axis
- $\|A_r - A_l\| =$ semi-minor axis
- $\delta_{rl} = \delta_l - \delta_r$
- $-\frac{1}{2}\delta_{rl} =$ position angle of MA
- $\delta_{rl} > 0$: MA rotated CW
- $\delta_{rl} = 0$: MA along x-axis
- $\delta_{rl} < 0$: MA rotated CCW


Polarization ellipse: circular 001




Polarization ellipse: circular 002




Polarization ellipse: circular 003




Polarization ellipse: circular 004




Polarization ellipse: circular 005




Polarization ellipse: circular 006




Polarization ellipse: circular 007




Polarization ellipse: circular 008




Polarization ellipse: circular 009




Polarization ellipse: circular 010




Polarization ellipse: circular 011




Polarization ellipse: circular 012




Polarization ellipse: circular 013




Polarization ellipse: circular 014




Polarization ellipse: circular 015




Polarization ellipse: circular 016




Polarization ellipse: circular 017




Polarization ellipse: circular 018




Polarization ellipse: circular 019




Polarization ellipse: circular 020




Polarization ellipse: circular 021




Polarization ellipse: circular 022




Polarization ellipse: circular 023




Polarization ellipse: circular 024




Polarization ellipse: circular 025




Polarization ellipse: circular 026




Polarization ellipse: circular 027




Polarization ellipse: circular 028



Polarization ellipse: circular 029



Polarization ellipse: circular 030



Relation between cartesian and circular

$$A_r = \frac{1}{2} \sqrt{A_x^2 + A_y^2 - 2A_x A_y \sin \delta_{xy}}$$

$$A_l = \frac{1}{2} \sqrt{A_x^2 + A_y^2 + 2A_x A_y \sin \delta_{xy}}$$

$$\tan \delta_{rl} = \frac{2A_x A_y \cos \delta_{xy}}{A_x^2 - A_y^2}$$

Stokes parameters

- Three parameters enough
- Same units is convenient
- George Stokes defined four parameters (1852)
- Chandrasekhar introduced them to astronomy (1946)

$$I = A_x^2 + A_y^2$$

$$Q = A_x^2 - A_y^2$$

$$U = 2A_x A_y \cos \delta_{xy}$$

$$V = -2A_x A_y \sin \delta_{xy}$$

$$I = A_r^2 + A_l^2$$

$$Q = 2A_r A_l \cos \delta_{rl}$$

$$U = -2A_r A_l \sin \delta_{rl}$$

$$V = A_r^2 - A_l^2$$

- **Monochromatic** wave 100% polarized:

$$I^2 = Q^2 + U^2 + V^2$$

Stokes parameters

- Three parameters enough
- Same units is convenient
- George Stokes defined four parameters (1852) *ABCD*
- Chandrasekhar introduced them to astronomy (1946) *I_II_rUV*

$$I = A_x^2 + A_y^2$$

$$Q = A_x^2 - A_y^2$$

$$U = 2A_x A_y \cos \delta_{xy}$$

$$V = -2A_x A_y \sin \delta_{xy}$$

$$I = A_r^2 + A_l^2$$

$$Q = 2A_r A_l \cos \delta_{rl}$$


$$U = -2A_r A_l \sin \delta_{rl}$$

$$V = A_r^2 - A_l^2$$

- *Monochromatic* wave 100% polarized:

$$I^2 = Q^2 + U^2 + V^2$$

IAU conventions



Public angry letter from IAU (Dec 2015)



International Astronomical Union
Union Astronomique Internationale
81bis bd Arago, F - 75014 Paris, France ■ www.iau.org

Piero Benvenuti - General Secretary
e-mail: iau-general.secretary@iau.fr

To Whom It May Concern

The Issue

Scientists working on the polarization of [REDACTED] use a convention for the polarization angle (PA) which is opposite to the IAU approved standard. This may cause confusion and misunderstandings.

Background

The convention astronomers follow for the PA (Polarization Angle) goes back to the 19th century and it has been in use for observations going from radio to gamma rays: the PA increases counter-clockwise when looking at the source. This convention is consistent with the one used for the Position Angle and it has been enforced by the IAU with a Resolution by Commissions 25 and 40 at the IAU XXth General Assembly in Sydney in 1973 (see Transactions of the IAU, Vol. XVI, pg. 166).

Recently, the scientists investigating the polarization of [REDACTED] have unfortunately adopted the opposite convention (PA increasing clockwise when looking at the source). This corresponds to a change of sign of the U Stokes parameter and is causing confusion and misunderstandings, in particular in the case of polarization data coming from experiments and satellites which are used by [REDACTED] and by other astronomers.

Recommendation

The IAU recommends that all astronomers, including those [REDACTED] follow the IAU Resolution for the Polarization Angle in all their publications.

Paris, December 8th, 2015

Piero Benvenuti
IAU General Secretary

Pietro Libertini
President, Division B

Saul J. Adelman
President, Commission B6

Executive Committee

Officers

Prof. Silvia Torres-Peimbert (Mexico), President
Prof. Piero Benvenuti (Italy), General Secretary
Prof. Ewine van Dishoeck (The Netherlands), President Elect
Prof. Maria Teresa Lago (Portugal), Assistant General Secretary

Vice Presidents

Prof. Rens C. Kraai-Worteweg (South Africa)
Prof. Xiaowei Liu (China)
Prof. Ding-Pei Li (Israel)
Prof. Debra Elmegreen (USA)
Prof. Ajit Kembhavi (India)
Prof. Boris Shustov (Russian Federation)

ASTRON

Netherlands Institute for Radio Astronomy

Stokes parameters revisited

Quasi-monochromatic approximation

- Monochromatic radiation does not exist
- Finite bandwidth $\Delta\nu$; averaging time $\tau \gg \Delta\nu^{-1}$

$$I = \langle A_x^2 \rangle + \langle A_y^2 \rangle$$

$$Q = \langle A_x^2 \rangle - \langle A_y^2 \rangle$$

$$U = \langle 2A_x A_y \cos \delta_{xy} \rangle$$

$$V = \langle -2A_x A_y \sin \delta_{xy} \rangle$$

$$I = \langle A_r^2 \rangle + \langle A_l^2 \rangle$$

$$Q = \langle 2A_r A_l \cos \delta_{rl} \rangle$$

$$U = \langle -2A_r A_l \sin \delta_{rl} \rangle$$

$$V = \langle A_r^2 \rangle - \langle A_l^2 \rangle$$

$$I^2 \geq Q^2 + U^2 + V^2$$


- Fractional linear pol: $p = \sqrt{Q^2 + U^2}/I \leq 1$
- Fractional circular pol: $v = \|V\|/I \leq 1$

Polarization orientation and Stokes parameters


(Vector length $\propto$ flux, not E -field;

Unit circle represents total flux I)


Polarization	I (Jy)	Q (Jy)	U (Jy)	V (Jy)
	1	0	0	0
	1	-1	0	0
	1	0	+1	0
	1	+1	0	0
	1	$+\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	0
	1	0	0	$-\frac{1}{2}$



TEST: Mars with VLA at 23.4 GHz *Perley*




- This is **thermal** emission
- 1) Draw a map of the pol vectors
- 2) Why is it even polarized?



Wait for it. . .

Answer: Mars polarization vectors *Perley*



Outline

EM wave physics


Astrophysics

Polarized EM-waves

Interferometric polarimetry

Messy reality

Stokes visibilities



$$\mathcal{I}(u, v) = \mathcal{F}^+(I(l, m))$$

$$\mathcal{Q}(u, v) = \mathcal{F}^+(Q(l, m))$$

$$\mathcal{U}(u, v) = \mathcal{F}^+(U(l, m))$$

$$\mathcal{V}(u, v) = \mathcal{F}^+(V(l, m)),$$

where

$$\mathcal{F}^+(f) = \int_{lm} f e^{+2\pi i \nu (ul + vm)/c} dl dm$$

Stokes parameters in complex notation

Cartesian

$$E_x = \Re \left\{ A_x e^{2\pi i \nu t} \right\}$$

$$E_y = \Re \left\{ A_y e^{i\delta_{xy}} e^{2\pi i \nu t} \right\}$$

$$I = \langle A_x^2 \rangle + \langle A_y^2 \rangle$$

$$= \langle E_x E_x^* \rangle + \langle E_y E_y^* \rangle$$

$$Q = \langle A_x^2 \rangle - \langle A_y^2 \rangle$$

$$= \langle E_x E_x^* \rangle - \langle E_y E_y^* \rangle$$

$$U = \langle 2A_x A_y \cos \delta_{xy} \rangle$$

$$= \langle E_x E_y^* \rangle + \langle E_y E_x^* \rangle$$

$$V = \langle -2A_x A_y \sin \delta_{xy} \rangle$$

$$= -i \left(\langle E_x E_y^* \rangle - \langle E_y E_x^* \rangle \right)$$

Stokes parameters in complex notation

Circular

$$E_r = A_r e^{2\pi i \nu t}$$

$$E_l = A_l e^{-i\delta_{rl}} e^{-2\pi i \nu t}$$

$$I = \langle A_r^2 \rangle + \langle A_l^2 \rangle$$

$$= \langle E_r E_r^* \rangle + \langle E_l E_l^* \rangle$$

$$Q = \langle 2A_r A_l \cos \delta_{rl} \rangle$$

$$= \langle E_r E_l^* \rangle + \langle E_l E_r^* \rangle$$


$$U = \langle -2A_r A_l \sin \delta_{rl} \rangle$$

$$= i (\langle E_r E_l^* \rangle - \langle E_l E_r^* \rangle)$$


$$V = \langle A_r^2 \rangle - \langle A_l^2 \rangle$$

$$= \langle E_r E_r^* \rangle - \langle E_l E_l^* \rangle$$

Correlating interferometer



Stokes visibilities from correlator outputs



- From here on, $\langle \cdot \rangle$ is implied for correlator outputs.

Cartesian

$$\mathcal{I} = x_1 x_2^* + y_1 y_2^*$$

$$\mathcal{Q} = x_1 x_2^* - y_1 y_2^*$$

$$\mathcal{U} = x_1 y_2^* + y_1 x_2^*$$

$$\mathcal{V} = -i(x_1 y_2^* - y_1 x_2^*)$$

Circular


$$\mathcal{I} = r_1 r_2^* + l_1 l_2^*$$

$$\mathcal{Q} = r_1 l_2^* + l_1 r_2^*$$

$$\mathcal{U} = i(r_1 l_2^* - l_1 r_2^*)$$

$$\mathcal{V} = r_1 r_2^* - l_1 l_2^*$$

Coherency matrix



From here on, p and q designate either x and y , or r and l .

- Polarizers produce vector:

$$\mathbf{e}_i = \begin{pmatrix} p_i \\ q_i \end{pmatrix}$$

- Correlator multiplies:

$$\mathbf{E}_{ij} = \mathbf{e}_i \mathbf{e}_j^\dagger = \begin{pmatrix} p_i \\ q_i \end{pmatrix} \begin{pmatrix} p_j^* & q_j^* \end{pmatrix}$$

$$\mathbf{E}_{ij} = \begin{pmatrix} p_i p_j^* & p_i q_j^* \\ q_i p_j^* & q_i q_j^* \end{pmatrix}$$

- $\mathbf{E}_{ij}$ is the **coherency matrix**

Outline

EM wave physics


Astrophysics

Polarized EM-waves

Interferometric polarimetry

Messy reality

Jones calculus



Until now...

- Assumed all systems perfect

From now...

- Assume all systems linear:

$$\mathbf{e}'_i = \mathbf{J}_i \mathbf{e}_i$$

- $\mathbf{J}_i$ (2×2) is **Jones matrix**
- Cross correlation:


$$\mathbf{E}'_{ij} = \mathbf{e}'_i \mathbf{e}'_j{}^\dagger$$

$$\mathbf{E}'_{ij} = \mathbf{J}_i \mathbf{e}_i (\mathbf{J}_j \mathbf{e}_j)^\dagger$$

$$\mathbf{E}'_{ij} = \mathbf{J}_i \mathbf{e}_i \mathbf{e}_j^\dagger \mathbf{J}_j^\dagger$$

$$\mathbf{E}'_{ij} = \mathbf{J}_i \mathbf{E}_{ij} \mathbf{J}_j^\dagger$$

Measurement equation



- The measurement equation:

$$\mathbf{E}'_{ij} = \mathbf{J}_i \mathbf{E}_{ij} \mathbf{J}_j^\dagger$$

- Invertible!


$$\mathbf{E}_{ij} = \mathbf{J}_i^{-1} \mathbf{E}'_{ij} \mathbf{J}_j^{\dagger-1},$$

- where

$$\mathbf{J} = \text{RPDOWFT}$$

- ... riiiiight...

Measurement equation: examples



- Perfect instrument:

$$\mathbf{J} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$


- Time delay:

$$\mathbf{J} = \begin{pmatrix} e^{2\pi i \nu \tau_p} & 0 \\ 0 & e^{2\pi i \nu \tau_q} \end{pmatrix}$$

- Receiver gain:

$$\mathbf{J} = \begin{pmatrix} g_p & 0 \\ 0 & g_q \end{pmatrix}$$

Measurement equation: examples



- Polarization leakage:

$$\mathbf{J} = \begin{pmatrix} g_p & d_{q \rightarrow p} \\ d_{p \rightarrow q} & g_q \end{pmatrix}$$



- Parallactic angle or feed rotation
XY:

$$\mathbf{J} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

- Parallactic angle or feed rotation
RL:

$$\mathbf{J} = \begin{pmatrix} e^{+i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}$$

Ionospheric Faraday rotation



- Remember: $\Delta\chi = \chi_0 + \phi\lambda^2$
- Faraday depth

$$\phi = 0.812 \int_{\text{there}}^{\text{here}} n_e \mathbf{B} \cdot d\mathbf{l}$$

- ionosphere: plasma within Earth's magnetic field
- $\phi \approx -10 - +10 \text{ rad m}^{-2}$
- *Very* significant below 1 GHz
- Use TEC/IGRF models for correction, check with pulsar.


Take away this...

- Radio antennas are **fundamentally polarized**
- **Polarimetry required** for certain astrophysical observations
- Linear systems make for fairly straightforward calibration
- Understanding polarimetry **improves** your **unpolarized** calibration and **imaging**

Reading list


- Born & Wolf *Principles of optics*
- Thompson, Moran & Swenson *Interferometry and Synthesis in Radio Astronomy*
- Taylor, Carilli & Perley *Synthesis Imaging in Radio Astronomy II*
- Bracewell *The Fourier Transform & Its Applications*
- Hamaker, Bregman & Sault *Understanding radio polarimetry: paper I*(1996)
- Sault, Hamaker & Bregman *paper II*(1996)
- Hamaker & Bregman *paper III* (1996)
- Hamaker *paper IV* (2000)
- Hamaker *paper V* (2006)
- Brentjens & de Bruyn *Faraday rotation measure synthesis* (2005)

TEST: Mars with VLA at 23.4 GHz *Perley*




- This is **thermal** emission
- 1) Draw a map of the polarization vectors.
- 2) Why is it even polarized?

Answer: Mars polarization vectors *Perley*



Faraday rotation measure synthesis




$$P(\lambda_c^2) = \int_{\lambda^{2'}} \int_{\Omega} \int_{-\infty}^{\infty} W(\Omega, \lambda_c^2 + \lambda^{2'}) f(\Omega, \phi, \lambda_c^2 + \lambda^{2'}) e^{2i\phi(\lambda_c^2 + \lambda^{2'})} d\phi d\Omega d\lambda^{2'}$$


$$P(\lambda^2) = W(\lambda^2) \int_{-\infty}^{\infty} f(\phi) e^{2i\phi\lambda^2} d\phi$$

$$f(\phi) * R(\phi) = \int_{-\infty}^{\infty} P(\lambda^2) e^{-2i\phi\lambda^2} d\lambda^2$$


Short wavelength



Intermediate wavelength




Long wavelength




Fan region *G. Bernardi*

RM: $-2.500000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.450000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.400000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.350000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.300000e+01$




Fan region *G. Bernardi*

RM: $-2.250000e+01$




Fan region *G. Bernardi*

RM: $-2.200000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.150000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.100000\text{e}+01$




Fan region *G. Bernardi*

RM: $-2.050000e+01$




Fan region *G. Bernardi*

RM: $-2.000000e+01$




Fan region *G. Bernardi*

RM: $-1.950000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.900000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.850000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.800000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.750000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.700000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.650000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.600000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.550000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.500000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.450000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.400000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.350000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.300000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.250000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.200000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.150000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.100000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.050000\text{e}+01$




Fan region *G. Bernardi*

RM: $-1.000000\text{e}+01$




Fan region *G. Bernardi*

RM: -9.500000e+00




Fan region *G. Bernardi*

RM: -9.000000e+00




Fan region *G. Bernardi*

RM: -8.500000e+00




Fan region *G. Bernardi*

RM: $-8.000000\text{e}+00$




Fan region *G. Bernardi*

RM: -7.500000e+00




Fan region *G. Bernardi*

RM: $-7.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-6.500000\text{e}+00$




Fan region *G. Bernardi*

RM: $-6.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-5.500000\text{e}+00$




Fan region *G. Bernardi*

RM: $-5.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-4.500000\text{e}+00$




Fan region *G. Bernardi*

RM: $-4.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-3.500000\text{e}+00$




Fan region *G. Bernardi*

RM: $-3.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-2.500000\text{e}+00$




Fan region *G. Bernardi*

RM: $-2.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-1.500000\text{e}+00$




Fan region *G. Bernardi*

RM: $-1.000000\text{e}+00$




Fan region *G. Bernardi*

RM: $-5.000000\text{e-}01$




Fan region *G. Bernardi*

RM: 0.000000e+00




Fan region *G. Bernardi*

RM: 5.000000e-01




Fan region *G. Bernardi*

RM: 1.000000e+00




Fan region *G. Bernardi*

RM: 1.500000e+00




Fan region *G. Bernardi*

RM: 2.000000e+00




Fan region *G. Bernardi*

RM: 2.500000e+00




Fan region *G. Bernardi*

RM: 3.000000e+00




Fan region *G. Bernardi*

RM: 3.500000e+00




Fan region *G. Bernardi*

RM: 4.000000e+00




Fan region *G. Bernardi*

RM: 4.500000e+00




Fan region *G. Bernardi*

RM: 5.000000e+00




Fan region *G. Bernardi*

RM: 5.500000e+00




Fan region *G. Bernardi*

RM: 6.000000e+00




Fan region *G. Bernardi*

RM: 6.500000e+00




Fan region *G. Bernardi*

RM: 7.000000e+00




Fan region *G. Bernardi*

RM: 7.500000e+00




Fan region *G. Bernardi*

RM: 8.000000e+00




Fan region *G. Bernardi*

RM: 8.500000e+00




Fan region *G. Bernardi*

RM: 9.000000e+00




Fan region *G. Bernardi*

RM: 9.500000e+00




Fan region *G. Bernardi*

RM: 1.000000e+01




Fan region *G. Bernardi*

RM: 1.050000e+01




Fan region *G. Bernardi*

RM: 1.100000e+01




Fan region *G. Bernardi*

RM: 1.150000e+01




Fan region *G. Bernardi*

RM: 1.200000e+01




Fan region *G. Bernardi*

RM: 1.250000e+01




Fan region *G. Bernardi*

RM: 1.300000e+01




Fan region *G. Bernardi*

RM: 1.350000e+01




Fan region *G. Bernardi*

RM: 1.400000e+01




Fan region *G. Bernardi*

RM: 1.450000e+01




Fan region *G. Bernardi*

RM: 1.500000e+01




Fan region *G. Bernardi*

RM: 1.550000e+01




Fan region *G. Bernardi*

RM: 1.600000e+01




Fan region *G. Bernardi*

RM: 1.650000e+01




Fan region *G. Bernardi*

RM: 1.700000e+01




Fan region *G. Bernardi*

RM: 1.750000e+01




Fan region *G. Bernardi*

RM: 1.800000e+01




Fan region *G. Bernardi*

RM: 1.850000e+01




Fan region *G. Bernardi*

RM: 1.900000e+01




Fan region *G. Bernardi*

RM: 1.950000e+01




Fan region *G. Bernardi*

RM: 2.000000e+01




Fan region *G. Bernardi*

RM: 2.050000e+01




Fan region *G. Bernardi*

RM: 2.100000e+01




Fan region *G. Bernardi*

RM: 2.150000e+01




Fan region *G. Bernardi*

RM: 2.200000e+01




Fan region *G. Bernardi*

RM: 2.250000e+01




Fan region *G. Bernardi*

RM: 2.300000e+01




Fan region *G. Bernardi*

RM: 2.350000e+01




Fan region *G. Bernardi*

RM: 2.400000e+01



Fan region *G. Bernardi*

RM: 2.450000e+01



Fan region (combined RGB)

